




SHOW PIG STARTER

#50MG

PROFILE

SHOW•RITE® Show Pig Starter is a carefully formulated medicated complete feed for optimal growth and performance. SHOW•RITE® Show Pig Starter is pelleted and incorporates the latest nutritional knowledge with good animal husbandry practices for maximum performance.



Show-Rite®

ABOUT THE PRODUCT:

SHOW•RITE® Show Pig Starter T135 C400

A complete swine starter formulated to be fed to starting pigs from 26 to 40 days of age.

Pelleted Feed

SHOW•RITE® Show Pig Starter is pelleted to help promote better feed consumption.

Medicated with Tiamulin and Chlortetracycline

For the control of swine dysentery associated with *Brachyspira hyodysenteriae* susceptible to Tiamulin. Treatment of bacterial enteritis caused by *Escherichia coli* and *Salmonella choleraesuis* and bacterial pneumonia caused by *Pasteurella multocida* susceptible to Chlortetracycline.

High Levels of Lysine

High levels of lysine and other amino acids insure maximum lean muscle deposition.

High Levels of Fat

High levels of fat add bloom and condition to high lean show pigs.

Added Phytase

Phytase is added to increase the digestibility of Phytin-bound phosphorus in swine diets



*Formulated to incorporate
the newest ingredient
technology available!*

The Name Says it All!

Product Number

#50MG—Show Pig Starter T135 C400, 50 lb.

Specialty 90-118

-continued on reverse side

NUTRIENT ANALYSIS:

Crude Protein min.....	21.0%
Lysine, min.....	1.5%
Crude Fat, min.....	7.0%
Crude Fiber, max.....	3.5%
Calcium, min.....	0.6%
Calcium, max.....	1.1%
Phosphorus, min.....	0.7%
Salt, min.....	0.3%
Salt, max.....	0.8%
Selenium, min.....	0.30 ppm
Zinc, min.....	3,000 ppm
Phytase (Aspergillus oryzae), min.....	227 FYT/lb*

*One unit of phytase activity (FYT) is defined as the quantity of enzyme which liberates 1 micromol of inorganic phosphorus per minute.

INGREDIENTS:

Grain products, animal protein products, plant protein products, pork fat (preserved with BHA), roughage products, calcium carbonate, monocalcium phosphate, dicalcium phosphate, menadione dimethylpyrimidinol bisulfite (source of vitamin K activity), L-lysine, vitamin B12 supplement, salt, DL-methionine, propionic acid (a preservative), biotin, niacin supplement, riboflavin supplement, vitamin D supplement, calcium silicate, pantothenic acid, copper sulfate, vitamin E supplement, phosphoric acid, L-threonine, choline chloride, manganese sulfate, pyridoxine hydrochloride, thiamine mononitrate, vitamin A acetate, manganous oxide, zinc sulfate, ferrous sulfate, calcium iodate, zinc oxide, dried Bacillus subtilus fermentation product, tartaric acid, sodium silico aluminate, L-tryptophan, ethylenediamine dihydriodide, ascorbic acid, sodium selenite, potassium chloride, artificial flavor, chromium tripicolinate, soybean oil, citric acid, yucca schidegera extract, malic acid, dried Aspergillus oryzae fermentation product (phytase).

FEEDING DIRECTIONS:

Feed continuously as the sole ration for 14 days.

Feed as the sole ration to starting pigs from 26 to 40 days of age at the rate of 0.5 pound per 10 pounds body weight to provide 10 milligrams per pound of Chlortetracycline per pound of body weight per day.

Feeding Tips:

- Water - Pigs should have access to clean, fresh water at all times.
- Make feed changes gradually, over at least a 5 day time period.
- Never feed any feedstuffs that are moldy, musty or suspect in any way.

SHOW•RITE® Show Pig Starter T135 C400 is formulated specifically for swine and should not be fed to other species.

WARNING: This product, which contains added copper, should not be fed to sheep or related species that have a low tolerance level to copper.

WARNING: Withdraw medicated feed 2 days before slaughter. Do not feed to animals other than swine. Use as the only source of Tiamulin and Chlortetracycline. Swine being treated with Tiamulin should not have access to feeds containing polyether ionophores (e.g., Monensin, Lasalocid, Narasin, Salinomycin and Semduramicin) as adverse reactions may occur. If signs of toxicity occur, discontinue use of medicated feeds.

SPECIAL CONSIDERATION: Protect medicated feeds from excessive temperatures. Do not store at warm temperatures, 104° F (40° C), for more than 7 days.

LIMITATIONS FOR USE: Feed continuously as sole ration for 14 days. Not for use in swine weighing more than 250 lb. Feed approximately 400 g/ton Chlortetracycline, varying with bodyweight and feed consumption to provide 10 mg/lb of bodyweight per day.